

Radiation burn-through measurements to infer opacity at conditions close to the solar radiative zone-convective zone boundary

David Hoarty¹, John Morton¹, Jonathan Rougier¹, Michael Rubery², Damon Swatton¹, Kyle McLean³, Robert Heeter², Scott Richardson¹, Steven Rose³, Ted Perry⁴, Yekaterina Opachich², Bruce Remington²

¹*Atomic Weapons Establishment, UK*

²*Lawrence Livermore National Laboratory, USA*

³*Imperial College London, UK*

⁴*Los Alamos National Laboratory, USA*

Recent measurements at the Sandia National Laboratory of the x-ray transmission of iron plasma much higher than predicted by theory have cast doubt on modelling of iron x-ray radiative opacity at conditions close to the solar convective zone-radiative zone boundary. An increased radiative opacity of the solar mixture, in particular iron, is a possible explanation for the disagreement in the position of the solar convection zone-radiative zone boundary as measured by helioseismology and predicted by modelling using the currently accepted elemental composition based on photosphere analysis. Here we present data from radiation burn-through experiments which do not support a large increase in the opacity of iron at conditions close to the base of the solar convection zone and provide a constraint on the possible values of both the mean opacity and the opacity in the x-ray range of the Sandia experiments.

Presenting Author Email Address: *david.hoarty@awe.co.uk*